

Contact Interactions and high Q^2 Events at HERA ¹

Dieter Zeppenfeld

Physics Department, University of Wisconsin, Madison, WI 53706

Abstract. Effective $eeqq$ contact interactions can enhance event rates in e^+p scattering at HERA at high Q^2 . Present constraints from atomic parity violation measurements and from Drell Yan events at the Tevatron are discussed.

Both the H1 [1] and ZEUS [2] experiments at HERA have observed an excess above SM expectations in $e^+p \rightarrow e^+jX$ events at $Q^2 > 15,000 \text{ GeV}^2$. Only the H1 data, however, are suggestive of an s -channel (leptoquark) resonance [3], and other possible sources of rate enhancements in high Q^2 DIS events need to be investigated.

A fairly model independent parameterization of possible new physics effects is provided by an effective Lagrangian of $eeqq$ contact interactions [4,5]: any exchange, in the s -, t - or u -channel, can be cast into this form provided the exchanged quanta are heavy compared to the center of mass energy of the scattering process. Following Ref. [5] the contact terms are parameterized as

$$L_{NC} = \sum_q \left[\eta_{LL} (\bar{e}_L \gamma_\mu e_L) (\bar{q}_L \gamma^\mu q_L) + \eta_{RR} (\bar{e}_R \gamma_\mu e_R) (\bar{q}_R \gamma^\mu q_R) \right. \\ \left. + \eta_{LR} (\bar{e}_L \gamma_\mu e_L) (\bar{q}_R \gamma^\mu q_R) + \eta_{RL} (\bar{e}_R \gamma_\mu e_R) (\bar{q}_L \gamma^\mu q_L) \right], \quad (1)$$

where the coefficients have dimension $(\text{TeV})^{-2}$ and are conventionally expressed as $\eta_{\alpha\beta}^{eq} = \pm 4\pi/\Lambda_{eq}^2$, with an effective mass scale of new physics, Λ_{eq} .

The contact terms (1) give additional contributions to the reduced amplitudes for $eq \rightarrow eq$, $\bar{q}q \rightarrow e^+e^-$, and $e^+e^- \rightarrow \bar{q}q$ subprocesses, beyond the SM photon- and Z -exchange terms:

$$M_{\alpha\beta}^{eq} = \frac{e^2 Q_e Q_q}{\hat{t}} + \frac{g_Z^2 (T_{e\alpha}^3 - s_w^2 Q_e)(T_{q\beta}^3 - s_w^2 Q_q)}{\hat{t} - m_Z^2} + \eta_{\alpha\beta}^{eq}. \quad (2)$$

Here Q_f and $T_{f\alpha}^3$ are charges and weak isospins, respectively, of the external fermions f_α , $g_Z = e/(\sin \theta_w \cos \theta_w)$, and $s_w = \sin \theta_w$. The Mandelstam invariant $\hat{t}$ is given by $\hat{t} = (E_{\text{c.m.}})^2$ for e^+e^- , by $\hat{t} = -Q^2 = -sxy$ for ep and by $\hat{t} = sx_1x_2$

¹⁾ Talk given at the *5th International Workshop on Deep Inelastic Scattering and QCD*, Chicago, Illinois, 14–18 April 1997.




FIGURE 1. Integrated cross sections versus a minimum Q^2 for $e^+p \rightarrow e^+X$ for the SM (solid curve) and two choices of contact interactions. The data points correspond to combined H1 [1] and ZEUS [2] measurements, assuming constant detection efficiencies of 80% and 81.5%, respectively. Errors are statistical only. From Ref. [5].

for $p\bar{p}$ collisions. Thus the presence of a contact interaction has interconnected implications for $ep \rightarrow ejX$, $p\bar{p} \rightarrow e^+e^-X$, and $e^+e^- \rightarrow \text{hadrons}$ and also for atomic physics parity violation experiments.

In ep scattering at high Q^2 valence quark contributions dominate and the cross sections are related to the amplitudes (2) and parton distribution functions $q(x, Q^2)$ as

$$\frac{d\sigma(e^+p)}{dx dy} \approx \frac{sx}{16\pi} \sum_{q=u,d} q(x, Q^2) \left[|M_{LR}^{eq}|^2 + |M_{RL}^{eq}|^2 + (1-y)^2 (|M_{LL}^{eq}|^2 + |M_{RR}^{eq}|^2) \right]. \quad (3)$$

From (3) we see that a high- y anomaly in e^+p requires an η_{LR}^{eq} or η_{RL}^{eq} amplitude. η_{LL}^{eq} or η_{RR}^{eq} amplitudes are suppressed by the $(1-y)^2$ factor in e^+p collisions but they would be enhanced compared to η_{LR} or η_{RL} terms in e^-p scattering. Because the d -parton density is severely suppressed in the large x region relative to the u -parton density, the eu contact interaction is needed to achieve an e^+p cross section enhancement.

As an illustration consider two scenarios, $\eta_{RL}^{eu} = \eta_{LR}^{eu} = 1.4 \text{ TeV}^{-2}$ and $\eta_{RL}^{eu} = -\eta_{LR}^{eu} = 2.6 \text{ TeV}^{-2}$, corresponding to scales $\Lambda_{eu} \approx 3 \text{ TeV}$ and $\Lambda_{eu} \approx 2.2 \text{ TeV}$, respectively. The resulting DIS cross sections, above a minimum Q^2 , are shown in Fig. 1. Both scenarios provide for a better description of the HERA data than the SM. However, their consistency with other data must be investigated.

The choice $\eta_{RL}^{eu} = -\eta_{LR}^{eu}$ is parity violating and modifies the SM prediction for the atomic physics parity violating parameter Q_W , given by

$$Q_W = Q_W^{\text{SM}} - \frac{1}{\sqrt{2}G_F} \left[(N + 2Z)\Delta\eta^{eu} + (2N + Z)\Delta\eta^{ed} \right]. \quad (4)$$

Here N is the number of neutrons, Z is the number of protons, and $\Delta\eta^{eq} = \eta_{RR}^{eq} - \eta_{LL}^{eq} + \eta_{RL}^{eq} - \eta_{LR}^{eq}$ is the coefficient of the VA combination of fermion currents which is probed. The $^{133}_{55}\text{Cs}$ measurements find $Q_W = -72.1 \pm 0.9$ [6] compared to a SM prediction of $Q_W^{\text{SM}} = -72.9$ [7] which yields

$$\Delta\eta^{eu} + 1.12\Delta\eta^{ed} = (-0.068 \pm 0.082) \text{ TeV}^{-2}. \quad (5)$$

The $\eta_{RL}^{eu} = -\eta_{LR}^{eu}$ scenario above is thus ruled out unless one introduces additional, cancelling terms such as $\eta_{LL}^{eu} = -\eta_{RR}^{eu} = \eta_{RL}^{eu}$.

Such additional terms would only marginally affect the HERA observables, but they would give rise to even larger deviations from the SM in $p\bar{p} \rightarrow \ell^+\ell^- X$ at the Tevatron, in particular at large $\ell^+\ell^-$ invariant mass, where contact interactions dominate the amplitudes and interference effects can be neglected. Recently, the CDF Collaboration has determined new preliminary limits on η_{LL}^{eq} contact terms corresponding to $\Lambda_{eq} > 3.4$ (2.4) TeV for $\eta_{LL}^{eu} < 0$ ($\eta_{LL}^{eu} > 0$) [8]. Comparison of the measured $\ell^+\ell^-$ invariant mass spectrum with predictions in the presence of contact terms indicate that the scenario $\eta_{RL}^{eu} = \eta_{LR}^{eu} = 1.4 \text{ TeV}^{-2}$ above is still (marginally) compatible with the present Tevatron data while scenarios such as $\eta_{RL}^{eu} = -\eta_{LR}^{eu} = 2.6 \text{ TeV}^{-2}$ are clearly ruled out. Present Tevatron Drell-Yan data probe contact terms at the same scale Λ_{eq} as HERA DIS data, but the two machines are complementary in their sensitivity to particular combinations of couplings.

ACKNOWLEDGMENTS

I would like to thank V. Barger, Kingman Cheung and K. Hagiwara for the enjoyable collaboration which lead to the results presented in this talk. This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

REFERENCES

1. H1 Collaboration, C. Adloff et al., report DESY 97-24 [hep-ex/9702012].
2. ZEUS Collaboration, J. Breitweg et al., report DESY 97-025 [hep-ex/9702015].
3. See contributions by J. Blümlein, G. Wang, J. Kalinowski, and S. Lola in these proceedings.

4. G. Altarelli et al., report CERN-TH/97-40 [hep-ph/9703276]; K. S. Babu et al., preprint IASSNS-HEP-97-04 [hep-ph/9703299]; N. Bartolomeo and M. Fabbrichesi, report SISSA-34-97-EP [hep-ph/9703375]; A. Nelson, report UW-PT-97-07 [hep-ph/9703379]; W. Buchmüller and D. Wyler, report DESY 97-066 [hep-ph/9704317]; N. G. Deshpande et al., report OITS-625 [hep-ph/9705236].
5. V. Barger et al., preprint MADPH-97-991 [hep-ph/9703311].
6. C.S. Wood et al., *Science* **275**, 1759 (1997).
7. Review of Particle Physics, *Phys. Rev.* **D54**, 1 (1996).
8. A. Bodek for the CDF Collaboration, preprint Fermilab-Conf-96/341-E (1996).